

Article type:
Original Research

Department of Dentistry, Al Hadi University, Iraq.

Corresponding author email address:
Shathaabdulwahid@huc.edu.iq



Article history:

Received 11 Jul 2025
Revised 20 Aug 2025
Accepted 25 Sep 2025
Published online 01 Nov 2025

How to cite this article:

Al Qaysi, SH. (2025). Salivary Stress Biomarkers During Academic Examinations in Second-Year Dental Students. International Journal of Body, Mind and Culture, 12(8), 43-48.



© 2025 the authors. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

Salivary Stress Biomarkers During Academic Examinations in Second-Year Dental Students

Shadha. Al Qaysi

ABSTRACT

Objective: Academic assessments are recognized as potent stressors for dental students. This study aimed to evaluate changes in salivary stress biomarkers—cortisol, α -amylase, and chromogranin A—associated with academic examinations in second-year dental students.

Methods and Materials: In this pre–post observational study, 40 second-year dental students (aged 20–22 years) from Al Hadi University College of Dentistry participated voluntarily. Unstimulated whole saliva was collected on two occasions: during a non-assessment period one week before examinations and during the assessment period at the end of exams. Samples were obtained by passive drooling for 5 minutes, coded to ensure confidentiality, and stored at -18°C until analysis. Salivary cortisol, α -amylase, and chromogranin A concentrations were measured using ELISA according to the manufacturers' instructions. Paired t-tests were used to compare biomarker levels between the two time points, with $p \leq 0.05$ considered statistically significant.

Findings: Significant differences were observed in salivary cortisol and chromogranin A between the non-assessment and assessment periods, with higher mean levels in the non-assessment period compared with the examination period. In contrast, salivary α -amylase showed no statistically significant change between the two time points. These results indicate that cortisol and chromogranin A are sensitive to changes across the academic assessment period, whereas α -amylase did not reflect these changes in this sample of dental students.

Conclusion: Academic examinations in second-year dental students are associated with measurable alterations in salivary cortisol and chromogranin A, supporting their use as non-invasive biomarkers of academic stress. Monitoring these biomarkers may help educators identify high-stress periods and design interventions to reduce stress and promote students' psychological well-being.

Keywords: Salivary biomarkers, academic stress, cortisol, dental students.

Introduction

Stress can be defined as a state of worry or mental tension due to difficult or unbearable situation. It can be defined also as a natural human response that allow persons to overcome life challenges and difficulties of life (Rocha et al., 2013).

Academic stress affects students' psychological and physiological well-being. Stress needs internal (psychological) balance of the body plus requirements of effort to adapted it (Crnković et al., 2018). Dental students often suffer from increased anxiety, depression and interpersonal sensitivity because of their studies

including theoretical and practical parts (Pani et al., 2011).

The stress has been shown to correlate with several salivary biomarkers. The choice of salivary biomarker to examine the body's response to academic stress depends on the type of stress studied. Chronic stress is associated with the activation of the hypothalamic-pituitary-adrenal (HPA) axis (measured by salivary cortisol), as well as with the depression of immune function (measured by salivary IgA and lysozyme). Acute stress is associated with activation of the sympatho-adrenomedullary system, which is reflected by salivary α -amylase and chromogranin A (Pani et al., 2011).

Saliva cortisol

Cortisol is thought to enter saliva by many ways independent of an active transport mechanism or by passive diffusion (Kirschbaum & Hellhammer, 1994). It reflects the secretory activity in hypothalamic-pituitary-adrenal (HPA) axis. Its level in saliva is less than that in blood. Salivary Cortisol correlates closer with free physiologically active serum cortisol fraction than with total serum cortisol, which contains the physiologically inactive protein-bound cortisol fraction (Takatsuji et al., 2008). The best advantage of salivary over serum cortisol measurement is the fear from needle will minimize the stress which may bias the results (Meeran et al., 1993). Salivary cortisol secretion, like serum cortisol, displays marked diurnal rhythm, its lowest level during slow-wave nocturnal sleep, then an increase during late sleep and reached its peak after awakening. Levels then slop rapidly followed by a sustained gradual decrease for the rest of the day (Edwards et al., 2001).

Salivary α -amylase is one of main enzyme in humans and is released from the salivary glands in response to adrenergic stimulation. α -Amylase breaks down starch, first to oligosaccharides and then to maltose and glucose, by hydrolysing α -1,4 glucan bonds. Amylase is also present in the pancreas, fallopian tubes, lung, prostate and ovarian tissues (Koh & Koh, 2007).

It has an immunological function, protecting the oral cavity from microorganisms (Scannapieco et al., 1993). Salivary α -amylase has also been recognized as a marker that is sensitive to stimuli activating the sympathetic system (e.g., adrenaline) (Ali & Nater, 2020). In health, α -amylase levels are the lowest in early morning and the highest in late afternoon (Takai et al., 2004). In response to stress, the concentration of

α -amylase in saliva rises rapidly, which makes it one of important salivary biomarker of stress (O'Donnell et al., 2009), specially, during blood sampling (Strahler et al., 2010). Acute stress activates the axis of sympathetic nervous system, adrenal medulla, which is reflected not only in the concentration of salivary α -amylase, but also in the concentration of salivary chromogranin A (Chojnowska et al., 2021; Engeland et al., 2016).

Chromogranin A (CgA)

CgA is an acidic glycoprotein released from the adrenal medulla and sympathetic nerve endings that can be measured in the saliva (Nagasawa et al., 1998; NISHIKAWA et al., 1998).

Its secretion is increased from submandibular salivary gland in response to different stress factors e.g., public speaking (NAKANE et al., 1998), sports (Lee et al., 2006), noise (Miyakawa et al., 2006; Ng et al., 2003), passing an academic exam (Takatsuji et al., 2008) medical intervention (Obara & Iwama, 2005; Hua et al., 2014). or driving a car (Ng et al., 2003). However, there are no reports on the level of salivary chromogranin A in people diagnosed with depression or anxiety, but the main advantage of determining salivary CgA that CgA levels are not affected by the time of the day. Although its concentrations reaches the peak shortly after waking up, they drop rapidly (within only 1 h), and remained constant through the whole day (NAKANE et al., 1998).

Methods and Materials

Study design and setting

This investigation was designed as a pre-post observational study carried out at Al Hadi University College of Dentistry. The study aimed to evaluate changes in salivary stress biomarkers associated with academic examinations in second-year dental students.

Participants and sampling

The study sample consisted of 40 second-year dental students aged between 20 and 22 years, who were enrolled in the Bachelor of Dental Surgery program at Al Hadi University College of Dentistry. Students were invited to participate voluntarily after receiving an explanation of the study objectives and procedures, and written informed consent was obtained from all participants. Only students who were available for both saliva collection sessions were included in the final analysis.

Saliva collection procedure

Saliva samples were collected on two occasions from each participant: the first sample was obtained one week before the examination period (non-assessment condition), and the second sample was collected at the end of the examination period (assessment condition). On each occasion, students were instructed to rinse their mouth with water to remove food debris and then wait for a short period before sampling. Unstimulated whole saliva was then collected by allowing the saliva to pool in the mouth and gently drooling it into a sterile, labeled collection tube. The tubes were immediately placed in an ice box and subsequently stored at -18°C until analysis. According to the manufacturer's instructions, all samples were analyzed within 7 days of collection.

Findings and Results

Table 1.

Comparison of salivary chromogranin A levels before and after academic assessment (n = 40)

Time point	Mean $\pm$ SE (ng/mL)	t-value	p-value
Before exams	462.62 $\pm$ 9.64		
After exams	434.15 $\pm$ 7.01	23.684	0.019

Note: Paired t-test; SE = standard error. S: Significant ($p \leq 0.05$).

Table 1 shows the comparison of salivary chromogranin A levels before and after the academic assessment period. The mean chromogranin A level before the exams ($M = 462.62$, $SE = 9.64$) was higher than after the exams ($M = 434.15$, $SE = 7.01$). A paired t-test

Table 2.

Comparison of salivary amylase levels before and after academic assessment (n = 40)

Time point	Mean $\pm$ SE (U/L)	t-value	p-value
Before exams	14.24 $\pm$ 0.55		
After exams	13.10 $\pm$ 0.37	1.384	0.095

Note: Paired t-test; NS: Non-significant ($p > 0.05$); SE = standard error.

Table 2 presents the comparison of salivary amylase levels before and after the academic assessment period. The mean amylase level slightly decreased from before the exams ($M = 14.24$, $SE = 0.55$) to after the exams ($M = 13.10$, $SE = 0.37$). However, the paired t-test revealed

Biochemical analysis

Salivary concentrations of cortisol, α -amylase, and chromogranin A were determined using commercially available enzyme-linked immunosorbent assay (ELISA) kits, following the protocols provided by the manufacturers. All assays were performed in duplicate to improve reliability of measurements.

Statistical analysis

For each biomarker, values obtained before and during the examination period were compared using paired statistical tests (e.g. paired t-test), with the level of significance set at $p \leq 0.05$. Data analysis was performed using standard statistical software.

indicated that this difference between the two time points was statistically significant, $t(39) = 23.684$, $p = 0.019$, demonstrating a significant pre-post change in chromogranin A across the assessment period.

that this difference was not statistically significant, $t(39) = 1.384$, $p = 0.095$. These findings indicate that salivary amylase did not show a significant change between the non-assessment and assessment conditions in this sample.

Table 3. *Comparison of salivary cortisol levels before and after academic assessment (n = 40)*

Time point	Mean $\pm$ SE ($\mu\text{g/L}$)	t-value	p-value
Before exams	26.33 $\pm$ 0.56		
After exams	24.31 $\pm$ 0.71	1.785	0.027

Note: Paired t-test; S: Significant ($p \leq 0.05$); SE = standard error.

Table 3 summarizes the comparison of salivary cortisol levels measured before and after the academic assessment period. The mean cortisol level before the exams ($M = 26.33$, $SE = 0.56$) was higher than after the

Discussion and Conclusion

Biological indicators for stress reactions are valuable markers in psychophysiological research. Academic examination stress is reported to increase physiological and psychological measures of stress and to decrease immune functioning. Psychosocial stress is widely known to induce various adaptational responses of physiologic systems with particular increasing activities in the hypothalamus-pituitary-adrenal axis (HPA) as well as in the sympathetic-adrenal-medullary (SAM) system. Cortisol levels reflect the HPA activity whereas SAA and chromogranin A are said to reflect the SAM (Tammayan et al., 2021).

Estimation of salivary stress biomarkers was done during periods of assessment exams (stress period) and one week before exam. This study clearly shows that there is a significant difference in the level of cortisol and chromogranin A stress biomarkers during assessment and non-assessment periods, clearly implicating the higher level of stress during exams. Salivary cortisol levels were notably less during non assessment period and increased significantly during the assessment period ($p=0.000$). Cortisol is a major glucocorticoid in humans that reflects adrenocortical activity. Activation of the HPA and subsequent release of cortisol are major components of the physiological stress response.

Salivary cortisol accurately reflects serum cortisol, (Špiljak et al., 2024). Many studies have reported increased cortisol levels during stressful situations such as academic examinations, cardiac surgeries and dental treatment procedures (Inder et al., 2012; Pruessner et al., 1997; Tammayan et al., 2021).

Alpha amylase

exams ($M = 24.31$, $SE = 0.71$). Results of the paired t-test showed that this difference was statistically significant, $t(39) = 1.785$, $p = 0.027$, indicating a significant pre-post change in salivary cortisol associated with the examination period.

The salivary enzyme alpha-amylase has been proposed as a marker for stress-induced activity of the sympathetic nervous system.

It shows nonsignificant difference between two reading which is not agreed with Tammayan et al and Ouda et al (Tammayan et al., 2021; Ouda et al., 2016).

Application of noradrenaline or beta-adrenergic agonist can stimulate sAA release which acts as an indicator of sympathetic activity. A variety of studies on stress have consistently found increased levels of sAA in response to stress and it is also generally used as a biomarker of stress (Tammayan et al., 2021; Ali & Nater, 2020).

In our study shows no significant difference due to its low durability in response to stress since the time between the onset of mental stressor and the peak in its level is very quick -between 60 seconds to 180 seconds and has a rapid recovery (Obayashi, 2013). It may be related to chronic stress not acute one according to Vineetha et al (Vineetha et al., 2014).

Chromogranin A

Chromogranin A is increased in secretion after the exam because the stress is effected by sympathetic nervous system which is agreed with (Takatsuji et al., 2008; Tammayan et al., 2021).

Its level also is increased during venipuncture in hospitalize children which could be considered as a good stress maker according to Lee et al.

Acknowledgments

The authors express their gratitude and appreciation to all participants.

Declaration of Interest

The authors of this article declared no conflict of interest.

Ethical Considerations

The study protocol adhered to the principles outlined in the Helsinki Declaration, which provides guidelines for ethical research involving human participants. Ethical considerations in this study were that participation was entirely optional.

Transparency of Data

In accordance with the principles of transparency and open research, we declare that all data and materials used in this study are available upon request.

Funding

This research was carried out independently with personal funding and without the financial support of any governmental or private institution or organization.

Authors' Contributions

All authors equally contribute to this study.

References

- Ali, N., & Nater, U. M. (2020). Salivary alpha-amylase as a biomarker of stress in behavioral medicine. *International journal of behavioral medicine*, 27(3), 337-342. <https://doi.org/10.1007/s12529-019-09843-x>
- Chojnowska, S., Ptaszyńska-Sarosiek, I., Kępa, A., Knaś, M., & Waszkiewicz, N. (2021). Salivary biomarkers of stress, anxiety and depression. *Journal of clinical medicine*, 10(3), 517. <https://doi.org/10.3390/jcm10030517>
- Crnković, D., Peco, M., & Gelo, J. (2018). Correlation between salivary biochemical stress indicators and psychological indicators. *Acta Clinica Croatica*, 57(2.), 316-326. <https://doi.org/10.20471/acc.2018.57.02.13>
- Edwards, S., Clow, A., Evans, P., & Hucklebridge, F. (2001). Exploration of the awakening cortisol response in relation to diurnal cortisol secretory activity. *Life sciences*, 68(18), 2093-2103. [https://doi.org/10.1016/S0024-3205\(01\)00996-1](https://doi.org/10.1016/S0024-3205(01)00996-1)
- Engeland, C., Hugo, F., Hilgert, J., Nascimento, G., Junges, R., Lim, H.-J., Marucha, P., & Bosch, J. (2016). Psychological distress and salivary secretory immunity. *Brain, behavior, and immunity*, 52, 11-17. <https://doi.org/10.1016/j.bbi.2015.08.017>
- Hua, J., Le Scanff, C., Larue, J., José, F., Martin, J.-C., Devillers, L., & Filaire, E. (2014). Global stress response during a social stress test: impact of alexithymia and its subfactors. *Psychoneuroendocrinology*, 50, 53-61. <https://doi.org/10.1016/j.psyneuen.2014.08.003>
- Inder, W. J., Dimeski, G., & Russell, A. (2012). Measurement of salivary cortisol in 2012—laboratory techniques and clinical indications. *Clinical endocrinology*, 77(5), 645-651. <https://doi.org/10.1111/j.1365-2265.2012.04508.x>
- Kirschbaum, C., & Hellhammer, D. H. (1994). Salivary cortisol in psychoneuroendocrine research: recent developments and applications. *Psychoneuroendocrinology*, 19(4), 313-333. [https://doi.org/10.1016/0306-4530\(94\)90013-2](https://doi.org/10.1016/0306-4530(94)90013-2)
- Koh, D. S.-Q., & Koh, G. C.-H. (2007). The use of salivary biomarkers in occupational and environmental medicine. *Occupational and environmental medicine*, 64(3), 202-210. <https://doi.org/10.1136/oem.2006.026567>
- Lee, T., Shimizu, T., Iijima, M., Obinata, K., Yamashiro, Y., & Nagasawa, S. (2006). Evaluation of psychosomatic stress in children by measuring salivary chromogranin A. *Acta paediatrica*, 95(8), 935-939. <https://doi.org/10.1080/08035250500538940>
- Meeran, K., Hattersley, A., Mould, G., & Bloom, S. (1993). Venepuncture causes rapid rise in plasma ACTH. *International Journal of Clinical Practice*, 47(5), 246-247. <https://doi.org/10.1111/j.1742-1241.1993.tb09686.x>
- Miyakawa, M., Matsui, T., Kishikawa, H., Murayama, R., Uchiyama, I., Itoh, T., & Yoshida, T. (2006). Salivary chromogranin A as a measure of stress response to noise. *Noise and Health*, 8(32), 108-113. <https://doi.org/10.4103/1463-1741.33951>
- Nagasawa, S., NISHIKAWA, Y., LI, J., FUTAI, Y., KANNO, T., IGUCHI, K., MOCHIZUKI, T., HOSHINO, M., YANAIHARA, C., & YANAIHARA, N. (1998). Simple enzyme immunoassay for the measurement of immunoreactive chromogranin A in human plasma, urine and saliva. *Biomedical Research*, 19(6), 407-410. <https://doi.org/10.2220/biomedres.19.407>
- NAKANE, H., ASAMI, O., YAMADA, Y., HARADA, T., MATSUI, N., KANNO, T., & YANAIHARA, N. (1998). Salivary chromogranin A as an index of psychosomatic stress response. *Biomedical Research*, 19(6), 401-406. <https://doi.org/10.2220/biomedres.19.401>
- Ng, V., Koh, D., Mok, B. Y., Chia, S. E., & Lim, L. P. (2003). Salivary biomarkers associated with academic assessment stress among dental undergraduates. *Journal of Dental Education*, 67(10), 1091-1094. <https://doi.org/10.1002/j.0022-0337.2003.67.10.tb03701.x>
- NISHIKAWA, Y., LI, J., FUTAI, Y., YANAIHARA, N., IGUCHI, K., MOCHIZUKI, T., HOSHINO, M., & YANAIHARA, C. (1998). Region-specific radioimmunoassay for human chromogranin A. *Biomedical Research*, 19(4), 245-251. <https://doi.org/10.2220/biomedres.19.245>
- <https://doi.org/10.2220/biomedres.19.237>
- O'Donnell, K., Kammerer, M., O'Reilly, R., Taylor, A., & Glover, V. (2009). Salivary α -amylase stability, diurnal profile and lack of response to the cold hand test in young women. *Stress*, 12(6), 549-554. <https://doi.org/10.3109/10253890902822664>
- Obara, S., & Iwama, H. (2005). Assessment of psychological tension after premedication by measurement of salivary chromogranin A. *Journal of clinical anesthesia*, 17(7), 554-557. <https://doi.org/10.1016/j.jclinane.2005.08.001>
- Obayashi, K. (2013). Salivary mental stress proteins. *Clinica chimica acta*, 425, 196-201. <https://doi.org/10.1016/j.cca.2013.07.028>
- Ouda, S., Alaki, S., Safi, M., Nadhreen, A., & Johani, K. (2016). Salivary stress biomarkers—Are they predictors of academic assessment exams stress. *J. Clin. Exp. Pathol*, 15, 276-279. <https://www.semanticscholar.org/paper/Salivary-Stress-Biomarkers-Are-They-Predictors-of-Ouda-Alaki/12bfe2edbf04b3b596f59a2cfaaa1b5b6ed82d8>

- Pani, S. C., Al Askar, A. M., Al Mohrij, S. I., & Al Ohali, T. A. (2011). Evaluation of stress in final-year Saudi dental students using salivary cortisol as a biomarker. *Journal of Dental Education*, 75(3), 377-384. <https://doi.org/10.1002/j.0022-0337.2011.75.3.tb05051.x>
- Pruessner, J. C., Wolf, O. T., Hellhammer, D. H., Buske-Kirschbaum, A., Von Auer, K., Jobst, S., Kaspers, F., & Kirschbaum, C. (1997). Free cortisol levels after awakening: a reliable biological marker for the assessment of adrenocortical activity. *Life sciences*, 61(26), 2539-2549. [https://doi.org/10.1016/S0024-3205\(97\)01008-4](https://doi.org/10.1016/S0024-3205(97)01008-4)
- Rocha, M. C. P. d., Martino, M. M. F. D., Grassi-Kassisse, D. M., & Souza, A. L. d. (2013). Stress among nurses: an examination of salivary cortisol levels on work and day off. *Revista da Escola de Enfermagem da USP*, 47(05), 1187-1194. <https://doi.org/10.1590/S0080-623420130000500025>
- Scannapieco, F. A., Torres, G., & Levine, M. J. (1993). Salivary α -amylase: role in dental plaque and caries formation. *Critical Reviews in Oral Biology & Medicine*, 4(3), 301-307. <https://doi.org/10.1177/10454411930040030701>
- Špiljak, B., Šimunović, L., Vilibić, M., Hanžek, M., Crnković, D., & Lugović-Mihić, L. (2024). Perceived stress, salivary cortisol, and temperament traits among students of dental medicine: a prospective and interventional study. *Behavioral sciences*, 14(4), 289. <https://doi.org/10.3390/bs14040289>
- Strahler, J., Mueller, A., Rosenloecher, F., Kirschbaum, C., & Rohleder, N. (2010). Salivary α -amylase stress reactivity across different age groups. *Psychophysiology*, 47(3), 587-595. <https://doi.org/10.1111/j.1469-8986.2009.00957.x>
- Takai, N., Yamaguchi, M., Aragaki, T., Eto, K., Uchihashi, K., & Nishikawa, Y. (2004). Effect of psychological stress on the salivary cortisol and amylase levels in healthy young adults. *Archives of oral biology*, 49(12), 963-968. <https://doi.org/10.1016/j.archoralbio.2004.06.007>
- Takatsuji, K., Sugimoto, Y., Ishizaki, S., Ozaki, Y., Matsuyama, E., & Yamaguchi, Y. (2008). The effects of examination stress on salivary cortisol, immunoglobulin A, and chromogranin A in nursing students. *Biomedical Research*, 29(4), 221-224. <https://doi.org/10.2220/biomedres.29.221>
- Tammayan, M., Jantaratnotai, N., & Pachimsawat, P. (2021). Differential responses of salivary cortisol, amylase, and chromogranin A to academic stress. *PLoS One*, 16(8), e0256172. <https://doi.org/10.1371/journal.pone.0256172>
- Vineetha, R., Pai, K.-M., Vengal, M., Gopalakrishna, K., & Narayanakurup, D. (2014). Usefulness of salivary alpha amylase as a biomarker of chronic stress and stress related oral mucosal changes—a pilot study. *Journal of clinical and experimental dentistry*, 6(2), e132. <https://doi.org/10.4317/jced.51355>